

Tsunami and active tectonics along the western margin of the Bering Sea - impact on the coastal zone environment and evolution

Tatiana Pinegina¹, Andrey Kozhurin²

¹ IVS FEB RAS, Institute of Volcanology and Seismology FEB RAS, Piip Boulevard 9, 683006 Petropavlovsk-Kamchatsky, Russia; email: tsunami@kscnet.ru

² Geological Institute, RAS, Pyzhevsky per. 7, 119017 Moscow, Russia

Over the last about 20 years, the Bering coast of Kamchatka was not considered as an area with a high level of earthquakes and tsunami risks, despite the 1969 Mw 7.7 tsunamigenic earthquake near the Ozernoi Peninsula. However, the 1991 Khailinskoe (Mw 6.6) earthquake in the southern part of the

Koryakia, and especially the 2006 Olytorskoe earthquake (Mw 7.6) had raised the public and scientific concern about the possibility of large (including tsunamigenic) earthquakes in this area.

The western Bering Sea overlies a tectonically complex region. The plate




Fig. 1: Recurrence interval of tsunami with runup >5 m at the different parts of the Bering Sea coast (based on tsunami deposits for the last ~2000 years).

in this region are not well established - geoscientists have proposed several different plate configurations. Multiplate models (Cook et al. 1986, Lander et al. 1994, Mackey et al. 1997, Apel et al. 2006) more easily explain the location and mechanisms of the 1969, 1991 and 2006 earthquakes that occurred on the inferred Okhotsk/ Bering/ and Bering/ North America plate boundaries. During 1998-2003 field seasons we studied paleotsunami deposits along the southwestern coasts of the Bering Sea. First, we examined geological evidence for the 1969 Ozernoy earthquake and tsunami (Bourgeois et al. 2006, Martin et al. 2008). Then, we used these data as well as data on other historical tsunamis, as a guide for analyzing more than 4000 years of paleoseismic record in the southwestern Bering Sea. In this area we have documented evidence for 12-15 tsunamis during about 4500 years. Based on tsunami runup (4-8 m) and tsunami inundation ($\leq 300-400$ m), we think that these events were produced by local earthquakes with $M_w \sim 7.5 \pm 0.5$. Possibly, by kinematics, they are underwater analogues of the Olytorskoe M_w 7.6 earthquake of April 20, 2006 in Koryakia (Pinegina, Konstantinova 2006, Pinegina, Kozhurin 2010).

In 2009-2010 we extended the paleoseismological investigation to the west and northwest coast of the Bering Sea. A number of active faults, deforming late Pleistocene-Holocene marine terraces were identified. These faults, probably, have a submarine continuation in the Bering Sea. Slip along these faults may generate tsunamigenic earthquakes. Based on our data, the recurrence interval of slips along a single active fault may be as long as several thousands to ~ 10 thousands of years. The recurrence interval of tsunami (with runup > 5 m) at the different parts of the Bering Sea coast vary, in average, from 125 years up to

~ 1000 years (Fig. 1). Historical data show that the tsunamis with sources situated along the Kurile-Kamchatka subduction zone do not influence the Bering Sea coast significantly. So, we suppose that most of the Bering Sea tsunamis come from local sources (Bourgeois et al. 2006). They may be generated by several zones located along the margins of the Komandorsky basin. The analysis of historical seismicity (1937-2010) clearly shows that possible tsunamigenic zones may be 1) at the western shelf of Komandorsky basin, its slope and foot, and 2) at the western end of the Aleutian Island Arc. To that, seismogenic and tsunamigenic zones may be 3) at the extension of active faults of the Stolbovskaya depression in the Pokaty canyon, and 4) at the continuation of active structures of Koriaksky highland in the Litke Strait. These last two zones are less clearly pronounced in the modern seismicity, and were identified mostly by our paleoseismological study. Still, we have no data, either historical or paleoseismological, about tsunami from local earthquakes in the Olytorskoy and Korf Bays. One known historical tsunami in these Bays with ~ 4 m runup was transoceanic (from Chili 1960). It is no question that such events if there were any, influenced geologic history of the coastal area. For example, the coastal coseismic deformations may cause rapid fluctuations in relative sea level and produce therefore great environmental changes in the lake's and lagoon's sedimentation as a whole. Tsunami waves can inject a large amount of allogenic material into the coastal area and modify it drastically. Study of earthquakes- and tsunami-related imprints on coastal sequences should give us possibility to better understand the interaction between onshore and offshore processes in context of reconstruction of the paleoenvironmental conditions and their evolution.

References

- Apel EV, Burgmann R, Steblou G, Vasilenko N, King R, Prytkov A (2006) Independent active microplate tectonics of northeast Asia from GPS velocities and block modeling. *Geophysical Research Letters* 33: L11303
- Bourgeois J, Pinegina T, Ponomareva V, Zaretskaia N (2006) Holocene tsunamis in the southwestern Bering Sea, Russian Far East, and their tectonic implications. *GSA bulletin* 118: 449-463

- Cook DB, Fujita K, McMullen C (1986) Present-day plate interactions in northeast Asia: North American, Eurasian, and Okhotsk plate. *J. Geodynamics* 6: 33-51
- Kozhurin AI (2004) Active faulting at the Eurasian, North American and Pacific plates junction. *Tectonophysics* 380: 273-285
- Kozhurin AI (2007) Active Faulting in the Kamchatsky Peninsula, Kamchatka-Aleutian Junction. In: Eichelberger J, Gordeev E, Izbekov P, Lees J (eds) *Volcanism and Subduction: The Kamchatka Region*. American Geophysical Union, Washington, DC: 263-282
- Lander AV, Bukchin BG, Droznin DV, Kiryushin AV (1994) Tectonic Position and focal parameters of the Hailinskoe (Koryaskoe) earthquake on 8 March, 1992: Is there a Bering Plate? *Geodynamics and Earthquake Forecasting: Computational Seismology* edition 26: 104-122
- Mackey KG, Fujita K, Gunbina L, Kovalev V, Imaev V, Kozmin B (1997) Seismicity of the Bering Strait region: evidence for a Bering block. *Geology* 25: 979-982
- Martin ME, Weiss R, Bourgeois J, Pinegina TK, Houston H, Titov VV (2008) Combining constraints from tsunami modeling and sedimentology to untangle the 1969 Ozernoi and 1971 Kamchatskii tsunamis. *Geophysical Research Letters* 35: L01610
- Pinegina TK, Konstantinova TG (2006) Macroseismic observation of consequences from April 21, 2006 “Olytorskoe” earthquake. *Bulletin of Kamchatka regional association "Educational-scientific center"*. *Earth Sciences* 7: 169-173 (in Russian)
- Pinegina TK, Kozhurin AI (2010) A new data on Olytorsky earthquake fault (Mw 7.6, April 21, 2006, Koriakia, Russia). *Bulletin of Kamchatka regional association "Educational-scientific center"*. *Earth Sciences* 16: 231-241 (in Russian)
- Seliverstov NI (2009) Geodynamic zones at the junction between the Kuril-Kamchatka and Aleutian Island Arcs. *Petropavlovsk-Kamchatsky, KamGU Vitusa Beringa*: 191 (in Russian)